

THE CRAFT CIDER REVIVAL

~ Some Technical Considerations

Presentation to SWECA 28th February 2007

Andrew Lea

SOME THINGS TO THINK ABOUT

- Orcharding and fruit selection
- Full juice or high gravity fermentations
- Yeast and sulphiting
- Keeping
- Malo-lactic maturation
- Style of finished product

What is your overall USP?

How are you differentiated?

CRAFT CIDER IS NOW SPREADING

Cidermaking was once widespread over the whole of Southern England
There are signs that it may be returning eg Kent, Sussex and East Anglia
So regional styles may be back in favour eg higher acid /less tannic in the East

CHOICE OF CIDER FRUIT

The traditional classification
(Barker, LARS, 1905)

	Acid %	'Tannin' %
Sweet	< 0.45	< 0.2
Sharp	> 0.45	< 0.2
Bittersharp	> 0.45	> 0.2
Bittersweet	< 0.45	> 0.2
<i>Finished Cider</i>	<i>~ 0.45</i>	<i>~ 0.2</i>

CHOICE OF "VINTAGE QUALITY" FRUIT

Term devised by Hogg 1886

Adopted by Barker 1910 to embrace superior qualities that could not be determined by analysis

This is still true today!

“VINTAGE QUALITY” LIST (1988)**Sharps / Bittersharps**

Dymock Red
Kingston Black
Stoke Red
Foxwhelp
Browns Apple
Frederick
Backwell Red

Bittersweets

Ashton Brown Jersey
Harry Masters Jersey
Dabinett
Major
White Jersey
Yarlington Mill
Medaille d’Or

Pure Sweets


Northwood
Sweet Alford
Sweet Coppin

BLENDING OR SINGLE VARIETALS?

Blending before fermentation can ensure good pH control (< 3.8)

High pH (bittersweet) juices prone to infection

Single varieties may be sensorially unbalanced unless ameliorated with dilution or added acid

RELATIONSHIP BETWEEN pH AND TITRATABLE ACID IS NOT EXACT

Most bittersweet juices are > pH 3.8 or < 0.4% titratable acidity. Above pH 3.8 or lower than 0.4% acid is a tricky area for microbial safety.

CHOICE OF ORCHARD TYPE**A traditional orchard**

30 standard trees per acre
 Sheep or cattle graze underneath
 Higher landscape value

An intensive orchard

300 bush trees per acre
 Grass cover with herbicide strip
 No livestock

INFLUENCE ON CIDER QUALITY?

Standard trees (especially in old orchards) tend to lower nitrogen levels
 They tend to ferment more slowly
 This may be beneficial to ultimate cider quality

LARS Data Pot Grown Dabinett

	Fed	Unfed
Leaf N %	2.34	2.00
Leaf K %	1.4	1.2
Juice SG	1.057	1.057
Juice pH	4.25	4.25
Tannin %	0.30	0.35
Yield (kg)	33	21
Juice N (mg/100ml)	6.8	3.3
Days Fermenting	32	70

FRUIT STORAGE?

Not all apples are ripe when picked

Mid /late season apples mature when stored
 Starch converts to sugar so higher potential alcohol
 Volatile flavour develops

Practical considerations are important too
 e.g. protection from pests, mould

EFFECTS OF MILLING AND PRESSING

Most people press straight away after milling..... but cuvage (pulp storage) may have benefits

Tannin oxidation generates more soluble colour in the first couple of hours
 Tannin can be significantly lost overnight if too harsh
 These effects depend on pulp layer thickness (ease of access of air)

Soluble pectin increases overnight (useful for later keeving)
 Other flavour precursors are developed

FLAVOUR PRECURSOR DEVELOPMENT

Some flavours and precursors are bound as glucosides

These are hydrolysed when pulp stands, producing
eg phenylethanol (floral odour); 1,3 octanediol

The later addition of acetaldehyde to 1,3 octane-diol during fermentation gives dioxolanes with 'green' cidery aromas

CHAPTALISATION ?

Raising SG with sugar to *ca* 1.100 to ferment into high alcohol base cider

This is diluted with water for retail sale

Finished cider has 30-50% juice equivalent

Saves on tank capacity and juice costs

Produces a lighter style of mass-market cider

FULL JUICE CIDER

This is now a selling point for some producers e.g. Henneys, Aspall

The 3CCPA Producers Charter requires a minimum 85% juice content

A useful point of differentiation in the market

SEASONALITY

Mainstream cider is made from concentrate and sugar syrup on a 2 week cycle throughout the year.
Follows brewery practice.

Craft cider is made on an annual seasonal rotation. Follows winery practice.

- Press in autumn.
- Ferment in winter
- Cask or bottle in spring
- Consume in summer (from cask) or at any later time up to 5 years (from bottle)

Craft cider is essentially 'slow food' – another USP!

THE FERMENTING YEAST

Wild yeasts are everywhere - maybe up to 10^4 cells per gram *inside* the apple

The inoculum stays on cloths and equipment - apple juice will ferment spontaneously with great ease

An ecological succession of yeasts takes place during traditional cider fermentation

Kloeckera apiculata starts ; dies at 2% alcohol

Saccharomyces cerevisiae finishes

THIS IS WHAT HAPPENS IN FRANCE

Mixed microflora are still prized

Little SO₂ is used - natural succession

Kloeckera and *Candida* to start

Saccharomyces to finish

Very slow cool seasonal fermentations (4° C)

More complex flavour profile
Kloeckera dominates at low temperatures

PURE CULTURE FERMENTATION

Defined strains of wine yeasts - *Saccharomyces cerevisiae* and *S. bayanus*
have been available **dried** for about 25 years (previously as liquid slope cultures)
Pitch at $5 * 10^6$ per ml into sulphited juice
Fast fermentation to high alcohol (~2 weeks) - important with chaptalised musts and for year round fermentations
Require added nutrients to perform effectively
Dependable flocculation
Predictable but arguably bland results

YEAST AND THE CRAFT CIDERMAKER

Are commercial wine yeasts really appropriate?
To work well they need high nutrient input
They work faster than a seasonal product needs
They tend to ferment to absolute dryness (not clear why?)

Ability to ferment high gravities is irrelevant
They are too wine like?

WILD YEAST SUCCESSIONS

Uncontrolled wild yeast fermentations can give very unpredictable results
BUT
Use of SO₂ to control mould, bacteria and to direct the yeast succession has a long history
sulphur candles and “sweetening” of wooden casks

Controlled use of SO₂ in wild yeast cider fermentations was re-introduced by Beech and Burroughs at Long Ashton in the 1960's

FIRE AND BRIMSTONE!

“Lay Brimstone on a Rag, and by a Wire let it down into the Cider-Vessel, and there fire it; and when the Vessel is full of the Smoak, the Liquor speedily pour'd in, ferments the better”

[Dr Beale FRS in Evelyn's *Pomona* 1664]

BACK TO THE 60'S!!

The 1660's or the 1960's
Do we need cultured yeasts at all?
If speed is not important, make a virtue of slowness
Revisit Beale, Barker, Burroughs and Beech
Wild does not mean wanton!
A little measured SO₂ keeps things under control and allows beneficial yeasts to develop

THE ROLE OF SULPHUR DIOXIDE

Controls bacterial and fungal infection

Inhibits undesirable yeasts if added with regard to correct concentration for the pH

Is essential if a 'pure culture' yeast is added and is to establish dominance

Less than 1 ppm of 'molecular SO₂' is required to be effective (*equiv* up to 200 ppm total SO₂)

SULPHUR DIOXIDE CAN DO MORE....

Acts as an antioxidant /reducing agent (SO₂ to SO₄)

Blocks haze formation by nucleophilic addition to 'tannin' polyphenols, preventing polymerisation and haze formation

But cannot be added during active fermentation since it binds strongly to acetaldehyde

Acts as an antimicrobial / antioxidant in storage (50 ppm is typically added there irrespective of pH)

USING SULPHUR DIOXIDE BEFORE FERMENTATION

Added most conveniently as 10% metabisulphite solution (5% SO₂)

Binds to juice components especially from rotten fruit – hence total SO₂ always *exceeds* free SO₂


Its chemical equilibrium is very pH dependent (pH 3.8 is practical limit) – so dose also depends on pH

Active range is 0.5 - 1 ppm (mg/l)

To achieve 1 ppm active molecular SO₂, up to 200 ppm might need to be added at pH 3.8

Lag phase while *S. cerevisiae* multiplies may be 2 weeks or longer.

Warning - If too much SO₂ is added at a low pH, **all** yeasts may be killed – so follow the chart!



(Note 200 ppm total SO₂ is legal limit.)

A DIVERSION THROUGH KEEVING...

'Keeving' is the old English and French method
Still common practice in Devon pre-1939
Allows production of naturally sweet cider
The juice stands for several days in the cold
The 'flying lees' or *chapeau brun* rises spontaneously to the top
The clear juice below is pumped off to ferment very slowly for the next three months

SCIENCE EXPLAINS TRADITION!

Apple pectin is a highly methoxylated polygalacturonic acid which is slowly de-esterified by an endogenous esterase
The liberated polygalacturonate (pectate) anion combines with free cations in the juice, notably calcium, asparagine and thiamin
This calcium pectate gel rises to the top buoyed up by CO₂ from the yeast

THE RESULT...

Clear juice depleted in amino nitrogen and thiamin (vitamin B1) - which bind to the pectate gel
Hence yeast nutrient and vitamin level is very low
So the fermentation is very slow and traditionally can take 3 - 4 months in the winter
Repeated racking stops fermentation
Cider may be filtered and bottled naturally sweet at SG 1.010 – 1.025
Slight carbonation develops in the bottle
Disappeared commercially post-1945 in UK

KEEVING IS OBSOLETE(?) IN ENGLAND


But is the norm in France as 'dépécation', where they now use:
Added commercial PME (*Rapidase CME*)
Added calcium (as the chloride)
Nitrogen bubble flotation in larger factories
Fermentation at ca 4° C
Frequent centrifugation to remove the yeast crop
This produces a naturally sweet cider of full flavour and low alcohol (a craft cidemaker's USP?)
There is a revival of interest in the UK and USA
Keeving 'kit' (chloride and enzyme) can be bought from Standa-Industrie in Caen

CIDER MATURATION

Traditional ciders finish fermenting in the spring
As the weather warms up and the trees bloom, the cider starts to 'work' again and becomes less harsh in flavour
Evidently the cider and the trees are somehow in sympathy!

THIS IS THE MALO-LACTIC CHANGE

Malic is the principal acid in apple. It is decarboxylated by lactic acid bacteria e.g. *Leuconostoc oenos* to give CO₂ and lactic acid



Hence the acidity can fall by 50% and the cider becomes slightly carbonated

Other interesting flavour changes occur too

e.g. Bacterial conversion of hydroxy-cinnamic acids
to smaller simple spicy phenols to give the “bittersweet aroma”
Likewise butteriness in Chardonnay

WOODEN VATS

Old oak vats largely provide an inoculum for malo-lactic organisms...
...unlike wine where oak flavours are transferred from new barrels
Wood can be a USP- but also a false friend!

Not all malo-lactic bacteria are good. Some may cause
Ropiness
Mousiness
Excess acid loss
Danger of working at $\text{pH} > 3.8$ without SO_2

CONTROLLING MALO-LACTIC CHANGE

Promote

Old wooden casks
Higher pH
Stand on lees
No SO_2 on storage
Add a commercial bacterial culture

Inhibit

Use sterilised tanks
Keep pH down
Rack off lees
Add SO_2 for storage
Use lysozyme to destroy the bacteria

FINING AND FILTRATION

Some ciders will ‘drop bright’
If not, add

- Egg white
- Ox blood
- Gelatin (+/- bentonite)
- Chitosan

(all are positively charged and work by neutralisation of negative charged particles in juice)

Followed by
Sheet or powder filter or centrifuge
And finally submicron membranes - ‘The Magic of the Millipore’

Nowadays cross-flow ultrafiltration replaces fining?

CIDER STORAGE

“L’air est l’ennemi mortel du cidre” (as the French say)

Keep all vessels closed and full at all times. Use SO_2 and or CO_2 / nitrogen blankets

FINISHED CIDER - DRY OR SWEET ?

Dry cider does not appeal to all (punters talk dry but buy sweet!)
sugar / acid / tannin balance
SG 1.005 - 1.020 is preferred range

Added sugar risks refermentation unless pasteurised. Alternatives are
Saccharin (bitter aftertaste)
Acesulfame-K (aspartame isn't stable long-term)
Sucralose (clean aftertaste)
These alternatives lack 'body' compared to sugar

FINISHED CIDER - FIZZY OR FLAT ?

Carbonation lifts the flavour of a cider and provides greater perceived acidity
Typically 5 - 12 g/l (1 - 4 atmospheres at 10°C) is used for sparkle

Saturation solubility of CO₂ is 2.5 g/l at 10°C
A sub-saturation carbonation of 1 g/l is normal for many still white wines
This level enhances flavour balance and body without any evident bubbles
Provides some anti-microbial effect

FINISHED CIDER - BOTTLE OR CASK? (I)

Cider was first bottled in the 1640's in the Forest of Dean
The slight continued fermentation gave 'natural condition' and sparkle
Pre-dates Champagne by at least 50 years!
The gentry had specially- made glasses to drink from

BOTTLED CIDERS

Traditional

Took cider to new markets (Bulmers 1900)
Can keep (and mature?) for years in glass
Yeast was often present in finished product
Risk of exploding bottles at SG > 1.005 (unless slack corks are used as in French *cidre bouché* today)
Virtually died out in UK after WW2

Modern

Sweetened, carbonated and pasteurised for stability
Life in PET is quite short (oxygen in, CO₂ out!)
Some bottle conditioned ciders are creeping back
True "*méthode champenoise*" is also possible (no yeast in finished product)

FINISHED CIDER - BOTTLE OR CASK? (II)

Cider in traditional cask is difficult to keep well past early summer
It needs protection from airborne contamination e.g. *Acetobacter* and *Brettanomyces*
Cider mugs were popular for cask ciders in 18th and 19th century taverns

CASK CIDERS

Traditional

Has to be drunk quickly once broached
Prone to acetification and mousy off-flavours on exposure to air
Can be 're-conditioned' by adding sugar and yeast
Haze may be a positive USP

Modern

Filtered, carbonated, pasteurised and sweetened into kegs
Beer dispense systems
Good storage life
Collapsible bag-in-box now available if carbonation is not required

CIDER QUALITY DEPENDS UPON.....

Juice Composition

apple juice (source, blending etc.)

Fermentation Control

yeast management (sulphite, nutrients etc.)

Post fermentation handling

directing maturation
preventing oxidation

THE CHOICES ARE YOURS!